
## **Unit 5 Lesson 3: More Movement**

# 1 Moving a Graph (Warm up)

### **Student Task Statement**


How can we translate the graph of A to match one of the other graphs?




#### 2 New Hours for the Kitchen

#### **Student Task Statement**

Remember the bakery with the thermostat set to  $65^{\circ}$  F? At 5:00 a.m., the temperature in the kitchen rises to 85°F due to the ovens and other kitchen equipment being used until they are turned off at 10:00 a.m. When the owner decided to open 2 hours earlier, the baking schedule changed to match.






hours after midnight

hours after midnight

- 1. Andre thinks, "When the bakery starts baking 2 hours earlier, that means I need to subtract 2 from x and that G(x) = H(x-2)." How could you help Andre understand the error in his thinking?
- 2. The graph of F shows the temperatures after a change to the thermostat settings. What did the owner do?




hours after midnight


3. Write an expression for F in terms of the original baking schedule, H.

### **3 Thawing Meat**

#### **Student Task Statement**

A piece of meat is taken out of the freezer to thaw. The data points show its temperature M, in degrees Fahrenheit, t hours after it was taken out. The graph M = G(t), where  $G(t) = -62(0.85)^t$ , models the shape of the data but is in the wrong position.





Jada thinks changing the equation to  $J(t) = -62(0.85)^t + 75.1$  makes a good model for the data. Noah thinks  $N(t) = -62(0.85)^{(t+1)} + 68$  is better.

- 1. Without graphing, describe how Jada and Noah each transformed the graph of G to make their new functions to fit the data.
- 2. Use technology to graph the data, J and N, on the same axes.
- 3. Whose function do you think best fits the data? Be prepared to explain your reasoning.

## Images for Activity Synthesis

